



## Evaluation of marine phytoplankton toxicity by application of marine invertebrate bioassays

Eva Aylagas<sup>1</sup>, Iratxe Menchaca<sup>1</sup>, Aitor Laza-Martínez<sup>2</sup>, Sergio Seoane<sup>2</sup>, Javier Franco<sup>1</sup>

<sup>1</sup> AZTI-Tecnalia, Marine Research Division, Herrera Kaia, Portualdea, z/g, 20110 Pasaia, Spain.

<sup>2</sup> Department of Plant Biology and Ecology, University of the Basque Country (UPV/EHU), Bilbao, Spain.  
E-mail: [eylagas@azti.es](mailto:eylagas@azti.es)

**Summary:** The dinoflagellate *Alexandrium minutum* and the haptophyte *Prymnesium parvum* are well known for their toxin production and negative effects in marine coastal environments. *A. minutum* produces toxins which cause paralytic shellfish poisoning in humans and can affect copepods, shellfish and other marine organisms. Toxins of *P. parvum* are associated with massive fish mortalities resulting in negative impacts on the marine ecosystem and large economic losses in commercial aquaculture. The aim of this work is to improve our knowledge about the reliability of the use of marine invertebrate bioassays to detect microalgae toxicity, by performing: (i) a 24- to 48-h test with the brine shrimp *Artemia franciscana*; (ii) a 48-hour embryo-larval toxicity test with the sea urchin *Paracentrotus lividus*; and (iii) a 72-h test with the amphipod *Corophium multisetosum*. The results indicate that *A. franciscana* and *P. lividus* larvae are sensitive to the toxicity of *A. minutum* and *P. parvum*. LC<sub>50</sub> comparison analysis between the tested organisms reveals that *A. franciscana* is the most sensitive organism for *A. minutum*. These findings suggest that the use of different organizational biological level bioassays appears to be a suitable tool for *A. minutum* and *P. parvum* toxicity assessment.

**Keywords:** *Alexandrium minutum*; *Prymnesium parvum*; *Paracentrotus lividus*; *Artemia franciscana*; *Corophium multisetosum*; toxicity bioassays.

### Evaluación de la toxicidad de fitoplancton marino mediante la aplicación de bioensayos con invertebrados marinos

**Resumen:** Las microalgas *Alexandrium minutum* y *Prymnesium parvum* son bien conocidas por la producción de toxinas y sus efectos negativos en nuestras costas. *A. minutum* produce la toxina PSP (Paralytic Shellfish Poisoning) que afecta a humanos y una gran variedad de organismos marinos como copépodos y marisco. Las toxinas de *P. parvum* producen grandes mortalidades de peces ocasionando graves impactos negativos en el ecosistema marino e importantes pérdidas económicas en acuicultura. Este trabajo tiene como objetivo mejorar el conocimiento del uso de bioensayos con invertebrados marinos en la detección de la toxicidad de microalgas, mediante: (i) un test de 24-48 horas con *Artemia franciscana*; (ii) un test de 48 horas con embriones y larvas del erizo de mar *Paracentrotus lividus*; y (iii) un test de 72 horas con el anfípodo *Corophium multisetosum*. Los resultados muestran que las larvas de *A. franciscana* y *P. lividus* son sensibles a la toxicidad de *A. minutum* y *P. parvum*. Los análisis LC<sub>50</sub> revelan que *A. franciscana* es el organismo más sensible a *A. minutum*. Estos resultados indican que los bioensayos con diferentes niveles de organización biológica parecen ser una herramienta apropiada para la evaluación de la toxicidad de *A. minutum* y *P. parvum*.

**Palabras clave:** *Alexandrium minutum*; *Prymnesium parvum*; *Paracentrotus lividus*; *Artemia franciscana*; *Corophium multisetosum*; bioensayos de toxicidad.

**Citation/Como citar este artículo:** Aylagas E., Menchaca I., Laza-Martínez A., Seoane S., Franco J. 2014. Evaluation of marine phytoplankton toxicity by application of marine invertebrate bioassays. *Sci. Mar.* 78(2): 173-183. doi: <http://dx.doi.org/10.3989/scimar.03957.26C>

**Editor:** E. Garcés.

**Received:** September 12, 2013. **Accepted:** January 31, 2014. **Published:** May 26, 2014.

**Copyright:** © 2014 CSIC. This is an open-access article distributed under the Creative Commons Attribution-Non Commercial License (by-nc) Spain 3.0.

## INTRODUCTION

Over the last few decades, the incidence of toxic phytoplankton blooms has apparently been increasing in coastal waters worldwide (GEOHAB 2012). Toxic

blooms cause negative impacts and economic losses in many parts of the world and their consequences and mechanisms depend on the species involved. Broadly, there are four categories of deleterious effects: risks to human health; loss of natural or cultured seafood re-

sources; impairment of tourism and recreational activities; and damage to non-commercial marine resources and wildlife (GEOHAB 2001, Garcés and Camp 2011, Anderson et al. 2012). Several taxonomic groups of microalgae are known as harmful marine species: dinoflagellates, diatoms, raphidophyceans, cyanophytes and haptophytes (Moestrup 1994).

Among the genera responsible for harmful algal blooms (HABs), the genus *Alexandrium* is one of the most important, in terms of severity, diversity and distribution of bloom impacts (Anderson et al. 2012). *Alexandrium minutum* Halim is well known for the production of paralytic shellfish poisoning toxins and its presence can affect copepods, shellfish, and other marine organisms (Zhenxing et al. 2006). *A. minutum* was first described in the Mediterranean Sea forming a red water discoloration in Alexandria Harbour, Egypt (Halim 1960); it has since then been reported widely on the Mediterranean and Atlantic Spanish coast (Delgado et al. 1990, Franco et al. 1994, Frangópulos et al. 2000, Figueroa et al. 2007, Penna et al. 2008 and ICES 2011).

The haptophyte alga *Prymnesium parvum* N. Carter has been responsible for toxic events, with severe ecological impacts, in many parts of the world (Moestrup 1994). This organism produces a set of highly potent exotoxins, commonly called prymensins, that have been shown to have several biological effects, including ichthyotoxic, neurotoxic, cytotoxic, hepatotoxic and hemolytic activity towards a range of marine organisms (Igarashi et al. 1996, Manning and La Claire 2010). *P. parvum* was described for the first time in England (Carter 1937) and has been responsible for massive fish mortalities around the world (Granéli and Turner 2006, Johnsen et al. 2010).

Because of these deleterious effects, several methods for determining the toxicity of marine microalgae have been used to date. The most common is the mouse bioassay developed by the Japanese Ministry of Health and Welfare (Yasumoto et al. 1978). This method is characterized by being accurate and of short-term response, but the controversy in the use of mammals for bioassays has led to the search for and practice of new alternative bioassays (FAO 2005). In this research field, phytoplankton toxicity has been tested in marine organisms, including fish and invertebrates; the tests include mortality tests with crustaceans of the genus *Artemia* (Houdan et al. 2004, Zhenxing et al. 2006, Faimali et al. 2012). Moreover, toxicity of extracts from different diatoms has been evaluated in sea urchin embryos (Buttino et al. 1999, Hansen et al. 2003). It should be noted that the toxic capacity of some toxic microalgae is still unknown and that toxicity induction depending on the environmental conditions, mainly nutrient availability, has led to some controversy (Granéli and Johanson 2003, Rimmel et al. 2011).

Due to the high toxicity reported for *A. minutum* and other dinoflagellates, its toxicity level should be fully evaluated to prevent the risk of bloom episodes in coastal areas. In addition, the involvement of the genus *Prymnesium* in fish mortalities makes its toxicity evaluation important in order to ensure the good quality of areas with fish aquaculture and shellfish exploitation.

Toxicity may be variable between strains of potentially toxic species (e.g. Ichimi et al. 2002, Montoya et al. 2010, Varela et al. 2012). It is therefore necessary to evaluate the toxicity of local strains of known potentially toxic species, such as *Alexandrium minutum* and *Prymnesium parvum*.

In the present study, bioassays with three different marine invertebrates (with different levels of biological organization) were applied as a detection tool to assess the toxic capacity of these marine microalgae species. A set of ecotoxicological tests was carried out to investigate the effects of *A. minutum* and *P. parvum* using different treatments on larvae of the sea urchin *Paracentrotus lividus* (Lamarck 1816), the branchiopod *Artemia franciscana* (Leach 1819) and adults of the amphipod *Corophium multisetosum* (Stock 1952).

The branchiopod *A. franciscana* is used routinely as a test organism for screening in ecotoxicological studies (Sorgeloos et al. 1978). Its life cycle begins with the hatching of dormant cysts that, once rehydrated in salt water, start the development as nauplii larvae, which are the material for the bioassays. These crustaceans are characterized by their adaptability to a wide range of salinity and temperature conditions; their short life cycle; their high adaptability to adverse environmental conditions; their high fecundity; their bisexual/parthenogenetic reproduction strategy (with nauplii or cysts production); their small body size; and their adaptability to variable food resources. They are also non-selective filter feeders sensitive to toxic substances (Sarabia 2002, Faimali et al. 2012).

The sea urchin *Paracentrotus lividus* is also used widely for testing particular contaminants (Fernández 2002) and determining phytotoxicity (Buttino et al. 1999, Hansen et al. 2003, Privitera et al. 2012). *P. lividus* has a wide geographic distribution, is abundant, and is easily harvested and maintained in the laboratory (Garmendia et al. 2009). Moreover, obtaining gametes and in vitro fecundation do not involve difficulties, while the embryonic development is short, thus allowing viable larvae to be obtained in a short time (Garmendia et al. 2009).

Marine and estuarine amphipods are regarded as one of the most suitable model species for integrating the effects of multiple toxic compounds (USEPA 2001, OSPAR Commission 2005). However, *C. multisetosum* has not been used previously for determining microalgae toxicity.

The bioassays were applied with the aim of (i) evaluating the sensitivity of the abovementioned species to the toxicity of *A. minutum* and *P. parvum* in the presence and absence of cells; and (ii) comparing the sensitivity of the three organisms to the toxicity of the mentioned phytoplankton species.

## MATERIALS AND METHODS

### Algae species

#### *Strains origin*

Microalgae samples were collected in the spring and summer of 2004 in the Abra of Bilbao, a bay at the sea-




Fig. 1. – The Basque Coast, showing the sampling localities. Microalgae sampling points: A, *Alexandrium minutum* (x) and *Prymnesium parvum* (y); B, sea urchin *Paracentrotus lividus*; C, amphipod *Corophium multisetosum*.

ward end of the Nervión estuary, in the Bay of Biscay. The clonal strain Dn4EHU of *Alexandrium minutum* was isolated from a sample collected in May 2004 in the marina of Getxo (inner Abra of Bilbao) (Fig. 1A). For strain isolation, a single cell was captured with a glass capillary under an inverted microscope (Nikon Diaphot TMD). Before being inoculated into the culture plate, the cell was transferred to a sterile culture medium drop and captured again, to eliminate possible contaminants. For the taxonomic identification, cells were stained with calcofluor ( $0.15 \text{ mg mL}^{-1}$ ) and thecal plates were observed using a Leica<sup>TM</sup> DMRB light microscope.

The *Prymnesium parvum* strain (Hp13EHU) was isolated from a plankton sample collected in August 2004 in the outer Abra of Bilbao (Fig. 1A). In order to obtain unialgal cultures, serial dilution cultures were made from the field sample. Taxonomic identification was performed through cellular scale characterization, using transmission electron microscopy (Seoane et al. 2009).

#### Culture conditions

*A. minutum* and *P. parvum* were grown in 1 L Erlenmeyer flasks containing 0.5 L of culture medium and maintained at  $20^\circ\text{C}$ , at an approximate irradiance of  $100 \mu\text{mol m}^{-2} \text{ s}^{-1}$  in a 14:10 light:dark period. Cultures were grown in f/2-Si (Guillard and Ryther 1962) prepared with filtered and autoclave-sterilized seawater. To obtain the growth curve, fluorescence

was measured daily with a Turner Designs 10-100R fluorometer as a proxy of cell density. Toxicity tests were performed with both cultures (*A. minutum* and *P. parvum*) in their early stationary growth stage in order to obtain high start-point cell densities.

Before being used for toxicological testing, the cell density was determined with a Neubauer chamber. Cultures were diluted with marine filtered water ( $0.22 \mu\text{m}$ ) (MFW) to obtain different algal concentrations.

#### Test organisms

##### *Paracentrotus lividus*

Fourteen *P. lividus* individuals with a test diameter (excluding spines) greater than 40 mm were collected from a rocky shore located on the west coast of Donostia-San Sebastián (Fig. 1B) in May and June 2012. Specimens were transported alive in a container with seawater and processed immediately in the laboratory.

##### *Artemia franciscana*

*Artemia franciscana* tests were performed following the standard operational procedure (Artoxkit 1990). Cysts (MicroBioTests Inc., Nazareth, Belgium) were hatched in 10 mL of standard seawater at  $25^\circ\text{C}$  under continuous light for 30 hours. After that time, all larvae had moulted to instar II-III stages and the hatched nauplii were separated from remaining cysts using a Pasteur pipette.

### *Corophium multisetosum*

Six hundred specimens of *C. multisetosum* were collected from sediments of the Bidasoa estuary (Fig. 1C) in June 2012 during low tide and transported to the laboratory. Individuals were maintained for one week following the USEPA (2001) recommendations; by being placed in trays containing water with continuous aeration, at 20°C and a salinity of 25 with an increase of three units per day until it reached salinity of 35.

### Toxicity bioassays

The ecotoxicological tests were performed using different cell concentrations and two treatments: test organisms were exposed (i) to the microalgae culture (cells + medium) and (ii) to the medium devoid of cells after eliminating the cells by filtration in order to detect the toxicity caused by the excretion of the toxins.

The bioassays were performed on II-III stage nauplii larvae of *Artemia franciscana*, on newly-fertilized sea urchin embryos (*Paracentrotus lividus*), and on adults of the amphipod *Corophium multisetosum*.

In order to define the accurate microalgae concentration range of *Alexandrium minutum* and *Prymnesium parvum* for the bioassay with *P. lividus* and *A. franciscana*, an initial test was undertaken. The mortality observed in this initial experiment helped us to define a more precise concentration range for a definitive test (see Table 1 for concentration ranges).

#### *P. lividus*: toxicity bioassay

The procedure followed in this study to carry out the *Paracentrotus lividus* embryo-larval bioassay, was taken from the protocol of Garmendia et al. (2009). After equatorial dissection on the sea urchins (Fig. 2A), oocytes (500 µL) were transferred to a 100-mL flask filled previously with MFW, and a small quantity of sperms was then added. After 5-10 min, egg density and fecundation success were determined by taking four 20-µL replicates and counting the total eggs in the aforementioned volume and the number of eggs that contained the fertilization membrane (Fig. 2B). A minimum fecundation success of 90% was established as being indicative of the good quality of the model organisms (Garmendia et al. 2009).

The bioassay was based on the exposure of the newly-fertilized eggs to different concentrations of cultured and filtered microalgae for 48 hours (Table 1).

After fertilization, approximately 500 eggs were transferred to propylene glasses (6 replicates for each algae concentration and 11 for the control, following Garmendia et al. [2009] and Beiras et al. [2002]) containing 20 mL of toxic algae solution or MFW. The glasses were covered and stored in the incubation camera in darkness for 48 hours at 20°C. After incubation, one replicate per treatment was used to measure oxygen, salinity and temperature in order to ensure normal optimal conditions. In the remainder of the replicates, 40% formaldehyde was added to arrest larva development. Embryonic success was estimated by counting the normal pluteus larvae among the 100 first larvae observed. Larvae were considered normal when the four arms were correctly formed (Garmendia et al. 2009) (Fig. 2C). On the other hand, larvae were considered non-well-developed when no four arms were formed (Fig. 2D). To consider the test as valid, the embryo success in the control must be higher than 90% (Fernández 2002). A negative control was performed with the non-toxic algae *Isochrysis galbana* (strain Hp1EHU).

#### *A. franciscana*: toxicity bioassay

The toxicity tests were performed in 24-well polystyrene multiwell plates containing 1 mL of different cell concentrations (Table 1). Marine standard water was used as a control and *I. galbana* as a negative control. Ten nauplii larvae were transferred to each well and the multiwell plates were covered with a parafilm and stored at 20°C in darkness for 48 hours. After 24 and 48 hours, dead larvae were counted. Larvae were considered dead when no movement was detected after 10 seconds of observation and semi-dead when the movement had decreased compared with the non-affected larvae. To consider the test as valid, the mortality in the control must be lower than 10% (Artoxkit 1990). Dead larvae were observed under the microscope in order to determine the presence or absence of algae cells inside the digestive system.

#### *C. multisetosum*: toxicity bioassay

The experiment was based on the exposure of adult amphipods to different concentrations of *A. minutum* and *P. parvum* (Table 1). Ten amphipods were placed in the glass flasks containing 100 mL of different microalgae concentrations. Three replicates were prepared for each concentration. Flasks were stored in the incubation cabinet in darkness for 72 hours at 20°C. After that time,

Table 1. – *Alexandrium minutum* and *Prymnesium parvum* concentration ranges in the different bioassays.

Algae species	Test organism	Bioassay	Concentration (cells mL <sup>-1</sup> )
<i>Alexandrium minutum</i>	<i>P. lividus</i>	Initial test	0; 100; 500; 1000; 5000; 50000; 100000
		Definitive test	0; 50000; 60000; 75000; 90000; 100000
	<i>Artemia</i> sp.	Initial test	0; 100; 500; 1000; 5000; 50000; 100000
		Definitive test	0; 1000; 5000; 10000; 50000; 100000
	<i>C. multisetosum</i>	-	0; 40000; 60000; 75000; 100000
<i>Prymnesium parvum</i>	<i>P. lividus</i>	Initial test	0; 1000; 5000; 10000; 100000; 230000
		Definitive test	0; 100000; 150000; 180000; 200000; 250000; 300000
	<i>Artemia</i> sp.	Initial test	0; 1000; 5000; 10000; 100000; 230000
		Definitive test	0; 5000; 10000; 100000; 200000; 250000
	<i>C. multisetosum</i>	-	0; 50000; 100000; 150000; 205000




Fig. 2. – *Paracentrotus lividus* bioassay. A, equatorial dissection of the sea urchin before fertilization; B, oocyte with the fertilization membrane after fertilization; C, sea urchin larva completely developed with the four arms correctly separate; D, poorly developed sea urchin larva resulting after exposure to toxic sample.

amphipods were considered dead when no movements were registered within 60 seconds. To consider the test as valid, the mortality in the control must not be higher than 10% (Casado-Martínez et al. 2006).

#### Reference toxicant tests

The quality and sensitivity of the model organisms used in the different bioassays was evaluated by the use of reference substances. Sodium dodecyl sulphate (SDS) was used in a range from 2 to 8 mg L<sup>-1</sup> for *P. lividus* and ammonium in a range from 4 to 128 mg L<sup>-1</sup> for *C. multisetosum*.

#### Statistical analysis

The LC<sub>50</sub> and LC<sub>10</sub> values (the concentrations causing, respectively, 50 and 10% mortality or non-normal development), together with their 95% confidence intervals were calculated by fitting the survival data to a regression curve by a Probit analysis (Fernández 2002),

with the statistical package Statgraphics® Plus 5.0. The “goodness of fit” was checked using the  $\chi^2$  test. Following the methodology described by the United States Environmental Protection Agency (USEPA 2002), when data did not fit Probit requirements (normality and homoscedasticity) a graphical method was used to determine LC<sub>50</sub> value.

Following Pérez and Beiras (2010), the NOEC (no effective observed concentration) and LOEC (low effective concentration) values were calculated with the Kruskal-Wallis and the Mann-Whitney U non-parametric test. Differences were considered as significant when *p* was lower than 0.05.

#### RESULTS

The abiotic parameters in all microalgae culture flasks (oxygen, salinity and temperature) are reported in Table 2. The LC<sub>10</sub> and LC<sub>50</sub> values for SDS in *P. lividus* and the LC<sub>50</sub> value for NH<sub>4</sub><sup>+</sup> in *Corophium* are reported in Table 3.

Table 2. – Physico-chemical parameters measured in flasks of the different microalgae cultures.

Species	Experiment	Oxygen (%)	Salinity	Temperature (°C)
<i>Isochrysis galbana</i>	Negative control test <i>P. lividus</i>	119	-	-
<i>Isochrysis galbana</i>	Negative control test <i>Artemia</i>	113.5	33.1	23.6
<i>Alexandrium minutum</i>	Initial test	100.3	35.1	20.4
<i>Prymnesium parvum</i>	Initial test	101	35.4	21
<i>Alexandrium minutum</i>	Definitive test	92.7	34.6	21.2
<i>Prymnesium parvum</i>	Definitive test	92.8	35.4	21

Table 3. – Lethal concentration (mg L<sup>-1</sup>) for SDS in *Paracentrotus lividus* and for NH<sub>4</sub><sup>+</sup> in *Corophium multisetosum*, their respective confidence intervals (95%) and NOEC (mg L<sup>-1</sup>) and LOEC (mg L<sup>-1</sup>) values.

Toxicant reference	Test organism	Bioassay	LC <sub>10</sub> (mg L <sup>-1</sup> )	LC <sub>50</sub> (mg L <sup>-1</sup> )	NOEC (mg L <sup>-1</sup> )	LOEC (mg L <sup>-1</sup> )	Reference
SDS	<i>Paracentrotus lividus</i>	Initial test	3.02 (2.91 – 3.12)	4.05 (3.98 – 4.11)	2.1	3.15	Present study
		Definitive test	1.99 (1.91 – 2.06)	2.65 (2.61 – 2.70)	-	2.03	Present study
		-	-	3.95 (3.34 – 4.56)	-	-	Menchaca 2010
NH <sub>4</sub> <sup>+</sup>	<i>Corophium multisetosum</i>	-	-	25.47 (20.98 – 31.15)	8	16	Present study
		-	-	55 (26 – 114.9)	-	-	Pérez 2006

The LC<sub>50</sub>-SDS for *P. lividus* in the initial test was 4.05 mg L<sup>-1</sup>; it was 2.65 mg L<sup>-1</sup> in the definitive test. Furthermore, NOEC and LOEC values for the initial test were 2.1 and 3.15 mg L<sup>-1</sup>, respectively. LOEC for the definitive test was 2.03 mg L<sup>-1</sup> (Table 3). The LC<sub>50</sub>-NH<sub>4</sub><sup>+</sup> for *C. multisetosum* was 25.47 mg L<sup>-1</sup>. Moreover, NOEC and LOEC values were 8 and 16 mg L<sup>-1</sup>, respectively.

### *Paracentrotus lividus* bioassay

The highest concentration of the dinoflagellate, 100000 cell mL<sup>-1</sup>, caused the inhibition of the embryonic development in all of the sea urchin larvae while from the lowest concentration tested (100 cell mL<sup>-1</sup>, to 1000 cell mL<sup>-1</sup>) no significant differences from the control were detected (Fig. 3A). A toxic effect was observed at




Fig. 3. – Percentage of *Paracentrotus lividus* embryonic development inhibition as a function of algal cell concentration with standard error of the mean for *Alexandrium minutum* culture of the initial (A) and definitive (B) tests and for *Prymnesium parvum* culture (C) and filtrate (D). \*: Significant differences ( $p < 0.05$ ).

Table 4. – Lethal concentration (mg L<sup>-1</sup>) for *Alexandrium minutum* and *Prymnesium parvum* (LC<sub>10</sub> and LC<sub>50</sub>) and their respective confidence intervals (95%), NOEC (mg L<sup>-1</sup>) and LOEC (mg L<sup>-1</sup>) values.

Algae species	Test organism	Bioassay	LC10 (cell mL <sup>-1</sup> )	LC50 (cell mL <sup>-1</sup> )	NOEC (cell mL <sup>-1</sup> )	LOEC (cell mL <sup>-1</sup> )	Reference
<i>A. minutum</i>	<i>P. lividus</i>	Initial test	-	65381	1000	5000	Present study
		Definitive test	-	<50000	-	50000	Present study
	<i>Artemia</i> sp.	24 h	-	150629 (83816 - )	1000	5000	Present study
		48 h	645 (107.2-1659.6)	16218 (9332.6-30199.5)	1000	5000	Present study
	<i>C. multisetosum</i>	-	-	-	100000	-	Zhenxing et al. 2006 Present study
<i>P. parvum</i>	<i>P. lividus</i>	Initial test	-	-	100000	230000	Present study
		Definitive test	-	-	180000	200000	Present study
	<i>S. granularis</i>	-	-	3-400000	-	-	Hansen et al. 2003
		24 h	-	-	250000	-	Present study
	<i>Artemia</i> sp.	48 h	-	-	250000	-	Present study
		24 h	-	500000	-	-	Houdan et al. 2004
		48 h	-	<100000	-	-	Present study
	<i>C. multisetosum</i>	-	-	26500	-	-	Granéli and Johansson 2003
<i>C. multisetosum</i>	-	-	-	205000	-	Present study	

5000 cell mL<sup>-1</sup>, with approximately 15% of embryonic inhibition. At 50000 cell mL<sup>-1</sup> the inhibition reached 20%. Considering these observations, a new range was applied in a definitive test: from 50000, to 100000 cell mL<sup>-1</sup> (Table 1). The results showed a marked toxic effect at 50000 cell mL<sup>-1</sup>, while 100% of inhibition was achieved above 60000 cell mL<sup>-1</sup> (Fig. 3B). For the initial and definitive tests, LOEC values were 5000 and 50000 cell mL<sup>-1</sup>, respectively (Table 4). On the other hand, the filtrate of *A. minutum* at the maximum concentration did not show any toxic effect on the sea urchin's embryonic development (data not shown).

For the initial test of *Prymnesium parvum*, significant differences from the control in the development inhibition of sea urchin larvae ( $p < 0.05$ ) were observed in a concentration of 230000 cell mL<sup>-1</sup>. For the definitive test, the concentration range was established at between 100000 and 300000 cell mL<sup>-1</sup>. From 100000 to 180000 cell mL<sup>-1</sup>, development inhibition was similar to the control (Fig. 3C); at and above 200000 cell mL<sup>-1</sup>, significant differences from the control were observed ( $p < 0.05$ ). The highest development inhibition was reached at 300000 cell mL<sup>-1</sup> (22.8%), so LC<sub>50</sub> could not be calculated. NOEC and LOEC values were 180000 and 200000 cell mL<sup>-1</sup>, respectively (Table 4). Considering the filtrate of *P. parvum*, inhibition of more than 50% of embryo development was obtained at a concentration of 300000 cell mL<sup>-1</sup>, showing significant differences from the control sample ( $p < 0.05$ ) (Fig. 3D).

#### *Artemia franciscana* bioassay

The mortality of the 15% of nauplii population was induced at a concentration of 5000 cells mL<sup>-1</sup> of the dinoflagellate *A. minutum* after 24 hours of exposure; however, 50% of mortality was not achieved with the initial concentrations. Higher cell densities were therefore used to perform the definitive bioassay (Table 1). In addition, the exposure time was increased from 24 to 48 hours. Incubation with the dinoflagellate significantly affected nauplii mortality during both exposure periods ( $p < 0.05$ ). After 24 hours of exposure, significant differences from the control were observed from




Fig. 4. – Mortality after 24 h (dark grey bar) and 48 h (light grey bar) of nauplii larvae of *Artemia franciscana* exposed to *Alexandrium minutum* culture. Two controls were used: standard seawater (0.22 µm) and *Isochrysis galbana* as a negative control. \* significant differences ( $p < 0.05$ ).

5000 cell mL<sup>-1</sup> (19 % of mortality) ( $p < 0.05$ ), increasing with the concentration until 45% mortality at 100000 cell mL<sup>-1</sup> (Fig. 4). After 48 hours, at 50000 and 100000 cell mL<sup>-1</sup>, the mortality in the nauplii population was 58% and 81%, respectively. The effect of exposure time can be observed with LC<sub>50</sub> values at 24 and 48 h: 150629 and 16218 cell mL<sup>-1</sup>, respectively (Table 4). In addition, the minimum concentration with significant mortality was 5000 cell mL<sup>-1</sup> at 24 and 48 hours of exposure. Related to the filtrate of *A. minutum*, the maximum algae concentration (100000 cell mL<sup>-1</sup>) showed no significant differences from the control sample ( $p > 0.05$ ) (data not shown). On the other hand, *A. franciscana* nauplii survival was not affected by the presence of *Prymnesium parvum* cells or filtrate at maximum exposure concentration, i.e. 250000 cell mL<sup>-1</sup> ( $p > 0.05$ ) (data not shown). However, a sub-lethal effect was detected along the concentration gradient, i.e. the movement of the nauplii was reduced at maximum exposure concentration (250000 cell mL<sup>-1</sup>), without reaching a mortality effect.

Due to logistic difficulties, the presence of cell content inside the alimentary canal of *A. franciscana* could not be identified.




Fig. 5. – Mortality after 72 h of *Corophium multisetosum* exposed to culture of *Alexandrium minutum* (left) and *Prymnesium parvum* (right). No significant differences in mortality between control and samples were observed.

### *Corophium multisetosum* bioassay

In relation to *C. multisetosum* mortality results, after 72 hours of exposure to both microalgae, significant differences from the control were not observed for any concentration ( $p > 0.05$ ). However, an increase in mortality was observed throughout the concentration range of *Alexandrium minutum* (Fig. 5).

### DISCUSSION

Lethal concentrations ( $\text{mg L}^{-1}$ ) for SDS in *P. lividus* and for  $\text{NH}_4^+$  in *C. multisetosum*, as calculated in this study, lie within the range of those obtained previously by Menchaca (2010) and Pérez (2006), respectively (Table 3). Our results have therefore confirmed the good quality of the model organisms (differences in the quality of gonad material were observed in the two different periods where sea urchins were collected, as the  $\text{LC}_{50}$  value for SDS in *P. lividus* obtained in May (the initial test) was higher than the value obtained in June (the definitive test). This observation is in accordance with Garmendia et al. (2010), who also detected variability in the quality of the gonad material throughout the year. Therefore, this assumption must be considered when interpreting the differences found in results of the two sets of trials performed with *P. lividus* exposed to *A. minutum*.

The dinoflagellate *Alexandrium minutum* was toxic for *A. franciscana* nauplii at  $5000 \text{ cell mL}^{-1}$ , while the  $\text{LC}_{50}$  value increased with exposure time. The 48-hour exposure proved to be much more harmful, indicating that both concentration and exposure time contributed to the toxic response in the nauplii population. The toxic effect of *A. minutum* on *A. franciscana* nauplii survival has rarely been examined. Our results agree with those of Zhenxing et al. (2006), who found that four strains of *Alexandrium* spp. were lethal for *A. salina* at a density of  $2000 \text{ cell mL}^{-1}$ . Regarding the filtrate of *A. minutum*, the maximum concentration ( $100000 \text{ cell mL}^{-1}$ ) showed no toxic effect in the *A. franciscana* nauplii, which is explained by the fact that their toxicity is linked to the intact cell, as has been reported for *Ostreopsis cf. ovata* (Faimali et al. 2012)

In contrast, *A. franciscana* nauplii survival was not affected by the presence of *P. parvum* culture or filtrate at the maximum exposure concentration ( $250000 \text{ cell mL}^{-1}$ ). These results contrast with those of Houdan et al. (2004), who found an  $\text{LC}_{50}$  value of a *P. parvum* culture (in the stationary phase) lower than  $100000 \text{ cell mL}^{-1}$  for *A. salina*. Some authors have suggested that this sensitivity disagreement could be related to differences in microalgae culture conditions (Granéli and Johanson 2003). According to the aforementioned authors, excretion of toxic metabolites might be a general mechanism among toxic haptophytes when exposed to stress events, e.g. nutrient-deficient conditions. However, recent studies hypothesize that the majority of *Prymnesium* toxicity is cellular-based rather than related to exotoxins production (Rommel et al. 2011). Further, reduction of *A. franciscana* movement with exposure time, as shown in the present study at the maximum exposure concentration of *P. parvum*, is in accordance with the results of Rommel and Hambright (2012). These investigators observed that *Prymnesium* cells were attached to most exterior surfaces, to fish gills and to feeding and respiratory appendages in zooplankton, which contributed to the movement depletion of individuals. Moreover, they found that the number of attached cells increased with exposure time and density of *Prymnesium*.

The toxicity of *A. minutum* and *P. parvum* had not been evaluated previously using sea urchin embryo bioassays. However, a dose-dependent toxic effect of several marine microalgae on sea urchin cell division has been recorded in several studies (Paul and Fenical 1984, Lemée et al. 1993, Pedrotti et al. 1996, Buttino et al. 1999). In the present study, *A. minutum* was toxic to sea urchin embryos at  $5000 \text{ cell mL}^{-1}$  (LOEC value). These results are in accordance with those of Privitera et al. (2012), who demonstrated the toxic effect in competent sea urchin larvae when exposed to the dinoflagellate *Ostreopsis cf. ovata*. On the other hand, filtrate of the dinoflagellate showed no toxic effect on the development of the sea urchin. This scenario is also in agreement with Faimali et al. (2012), who demonstrated that dinoflagellates do not show significant toxicity with growth medium devoid of cells.

By contrast, the culture and filtrate of *P. parvum* significantly inhibited embryo development at 200000 and 300000 cell mL<sup>-1</sup>. These effective concentrations are within the range of those calculated by Hansen et al. (2003), who reported a cell division reduction of newly-fertilizing eggs of the sea urchin *Sphaerechinus granularis* exposed to a haptophyte. However, Rimmel and Hambright (2012) suggested that the presence of toxins in the filtrate could be an artefact caused by the vacuum pressure used in the filtration process, which can cause cell disruption and the subsequent release of intracellular toxins. The result of the filtrate reported in the present study should therefore be considered carefully.

In the present study, adult amphipods were not sensitive to the toxicity of *A. minutum* and *P. parvum* at the concentrations tested. This finding contrasts with those of other authors who have routinely used adult zooplankton organisms (*Euterpina acutifrons* and *Acartia grani*) to evaluate the toxicity of *A. minutum* (Frangópulos et al. 2000, Costa and Fernández 2002). However, Romano et al. (2003) observed that secondary metabolites, such as unsaturated aldehydes produced by some algae during stress events, such as depredation (Pohnert 2000), were not toxic in adult copepods, while they inhibited the development of oocytes and embryos. Romano et al. (2003) hypothesized that this behaviour could be a defence mechanism of microalgae, designed to regulate the populations of herbivorous zooplankton.

Assuming similar test conditions (photoperiod, temperature, salinity and oxygen) for all the bioassays performed, among the tested organisms *A. franciscana* nauplii showed the highest sensitivity for *Alexandrium minutum* according to LC<sub>50</sub>-48h. These results are in agreement with those of Faimali et al. (2012) and Privitera et al. (2012), who suggested that the development stages could affect the sensitivity of the organisms, increasing the mortality with the ingestion rate. Moreover, Zhenxing et al. (2006) confirmed a decrease in the survival of *Artemia salina* nauplii with the feeding rates of *Alexandrium* spp. However, in the present study, owing to logistic difficulties, the presence of cell content inside the alimentary canal of *A. franciscana* could not be identified. As a result, it cannot be concluded that the higher sensitivity of *A. franciscana* nauplii is related to the ingestion of the microalgae. In the case of *P. parvum*, difficulties in the LC<sub>50</sub> calculation did not permit comparison of the sensitivity between the tested organisms.

The presence of toxic microalgae on the coasts of the Basque Country is monitored every year, within the context of the EU Water Framework Directive (WFD) (e.g. Borja et al. 2009). In 2008 the cell density of *Alexandrium minutum* was lower than 100 cell L<sup>-1</sup>, much less than the value considered as a limit for toxicity risk found in the present study. *Prymnesium parvum* has not been reported specifically in the WFD reports. The inverted light-microscope technique based upon fixed samples used in this monitoring programme for phytoplankton identification and counting does not usually allow species or even genus-level identification, of

small naked flagellates such as *Prymnesium* spp. The species level identification of prymnesiophyceans requires the analysis of the body scales through electron microscopy (Seoane et al. 2009) or molecular methods.

Regarding toxicity of *Alexandrium minutum*, it must be considered that the long culture maintenance of the dinoflagellate in the present work may have led to the decrease in toxin production. Martins et al. (2004) demonstrated that saxitoxin production is not a constitutive, stable characteristic of *Alexandrium*, but can be lost during routine culture maintenance over a prolonged interval.

## CONCLUSION

Only larvae of *Paracentrotus lividus* and *Artemia franciscana* have been demonstrated to be sensitive to the toxicity of the two marine microalgae *Alexandrium minutum* and *Prymnesium parvum*. The algae strains tested were isolated much in advance compared with the performed test. Therefore, the minimum toxic concentration values established in the present study for the model organisms cannot be considered as a real toxic value. We suggest performing the bioassays of the present study with the model organisms, using recent isolated *A. minutum* or *P. parvum* during toxic blooms and combining different parameters (nutrients, light and temperature) to study the potential toxicity in different conditions. Nevertheless, our results place *P. lividus* and *A. franciscana* bioassays as a viable alternative for the detection of *A. minutum* and *P. parvum* toxicity in the context of routine monitoring. Other standardized toxicological assays using more distant taxonomic groups are also available and have been applied extensively (Ribo and Kaiser 1987, Zon and Peterson 2005). Therefore, besides working on the use of the bioassays described in the present work, we also suggest the use of alternative methodologies in order to explore a wider range of tests for the detection of toxic events.

## ACKNOWLEDGEMENTS

This paper constitutes part of the master's thesis of E. Ayllagas. We wish to thank Professor Michael Collins from the School of Ocean and Earth Science of the University of Southampton (UK) and Dr. Ángel Borja from AZTI-Tecnalia (Spain) for kindly advising us on some details of the manuscript. This paper is contribution number 661 from AZTI-Tecnalia (Marine Research Unit).

## REFERENCES

- Anderson D.M., Alpermann T.J., Cembella A.D. et al. 2012. The globally distributed genus *Alexandrium*: multifaceted roles in marine ecosystems and impacts on human health. *Harmful Algae* 14: 10-35.  
<http://dx.doi.org/10.1016/j.hal.2011.10.012>
- Artokkit M. 1990. *Artemia* toxicity screening test for estuarine and marine waters. Standard operational procedure, Creasel, Deinze, Belgium.
- Beiras R. 2002. Comparison of methods to obtain a liquid phase in marine sediment toxicity bioassays with *Paracentrotus lividus*

- sea urchin embryos. Arch. Environ. Con. Tox. 42: 23-28.  
<http://dx.doi.org/10.1007/s002440010287>
- Borja A., Bald J., Franco J. et al. 2009. Using multiple ecosystem components, in assessing ecological status in Spanish (Basque Country) Atlantic marine waters. Mar. Pollut. Bull. 59: 54-64.  
<http://dx.doi.org/10.1016/j.marpolbul.2008.11.010>
- Buttino I., Miralto A., Ianora A. et al. 1999. Water-soluble extracts of the diatom *Thalassiosira rotula* induce aberrations in embryonic tubulin organisation of the sea urchin *Paracentrotus lividus*. Mar. Biol. 134: 147-154.  
<http://dx.doi.org/10.1007/s002270050533>
- Carter N. 1937. New or interesting algae from brackish water. Arch. Protistenkd. 90: 1-68.
- Casado-Martínez M.C., Beiras R., Belzunce M.J et al. 2006. Ejercicio interlaboratorio de bioensayos marinos para la evaluación de la calidad ambiental de sedimentos costeros. IV. Ensayo de toxicidad sobre sedimento con crustáceos anfípodos. Cienc. Mar. 32: 149-157.
- Costa R.M., Fernández F. 2002. Feeding and survival rates of the copepods *Euterpina acutifrons* Dana and *Acartia grani* Sars on the dinoflagellates *Alexandrium minutum* Balech and *Gyrodinium corsicum* Paulmier and the *Chryptophyta* *Rhodomonas baltica* Karsten. J. Exp. Mar. Biol. Ecol. 273: 131-142.  
[http://dx.doi.org/10.1016/S0022-0981\(02\)00132-6](http://dx.doi.org/10.1016/S0022-0981(02)00132-6)
- Delgado M., Estrada M.J., Camp J.V. et al. 1990. Development of a toxic *Alexandrium minutum* Halim (Dinophyceae) bloom in the harbour of Sant Carles de la Ràpita (Ebro Delta, northwestern Mediterranean). Sci. Mar. 54: 1-7.
- Faimali M., Giussani V., Piazza V. et al. 2012. Toxic effects of harmful benthic dinoflagellate *Ostreopsis ovata* on invertebrate and vertebrate marine organisms. Mar. Environ. Res. 76: 97-107.  
<http://dx.doi.org/10.1016/j.marenvres.2011.09.010>
- FAO 2005. Biotoxinas marinas. Estudio FAO: Alimentación y Nutrición. Organización de las Naciones Unidas para la Agricultura y la Alimentación. Roma, Italy, 278 pp.
- Fernández N. 2002. Evaluación biológica de la contaminación marina costera mediante bioensayos con embriones del erizo de mar *Paracentrotus lividus*. PhD thesis, University of Vigo, 211 pp.
- Figueroa R.I., Garcés E., Bravo I. 2007. Comparative study of the life cycles of *Alexandrium tamatum* and *Alexandrium minutum* (Gonyaulacales, Dinophyceae) in culture. J. Phycol. 43: 1039-1053.  
<http://dx.doi.org/10.1111/j.1529-8817.2007.00393.x>
- Franco J.M., Fernández P., Reguera B. 1994. Toxin profiles of natural populations and cultures of *Alexandrium minutum* Halim from Galician (Spain) coastal waters. J. Appl. Phycol. 6: 275-279.  
<http://dx.doi.org/10.1007/BF02181938>
- Frangópulos M., Guisande C., Maneiro I. et al. 2000. Short-term and long-term effects of the toxic dinoflagellate *Alexandrium minutum* on the copepod *Acartia clausi*. Mar. Ecol.-Prog. Ser. 203: 161-169.  
<http://dx.doi.org/10.3354/meps203161>
- Garcés E., Camp J. 2011. Habitat changes in the Mediterranean Sea and the consequences for Harmful Algal Blooms formation. In: N. Stambler (ed.), Life in the Mediterranean Sea: A look at habitat changes. Nova Science Publishers, Inc. New York, US, ISBN: 978-1-61209-644-5, pp. 519-541.
- Garmendia J.M., Menchaca I., Belzunce M.J. et al. 2009. Protocolo del test de toxicidad de sedimentos marinos con larvas del erizo de mar *Paracentrotus lividus* (Lamarck, 1816). RIM. 11-25.
- Garmendia J.M., Menchaca I., Belzunce M.J. et al. 2010. Seasonal variability in gonad development in the sea urchin (*Paracentrotus lividus*) on the Basque coast (southeastern Bay of Biscay). Mar. Pollut. Bull. 61: 259-266.  
<http://dx.doi.org/10.1016/j.marpolbul.2010.02.023>
- GEOHAB 2001. Global Ecology and Oceanography of Harmful Algal Blooms, Science Plan. P. Gilbert, G. Pitcher (eds), SCOR and IOC, Baltimore and Paris, 87 pp.
- GEOHAB 2012. Global Ecology and Oceanography of Harmful Algal Blooms, Science Plan. Berdalet E., Tester P., Zingone A. (eds), IOC of UNESCO and SCOR, Paris and Newark, 64 pp.
- Granéli E., Johansson N. 2003. Increase in the production of allelopathic substances by *Prymnesium parvum* cells grown under N- or P-deficient conditions. Harmful Algae 2: 135-145.  
[http://dx.doi.org/10.1016/S1568-9883\(03\)00006-4](http://dx.doi.org/10.1016/S1568-9883(03)00006-4)
- Granéli E., Turner J.T. (eds) 2006. Ecology of Harmful Algae. Ecol. Stud. 189, Springer. 270 pp.
- Guillard R.R.L., Ryther J.H. 1962. Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt, and *Detonula confervacea* (Cleve) Gran. Can. J. Microbiol. 8: 229-239.  
<http://dx.doi.org/10.1139/m62-029>
- Halim Y. 1960. *Alexandrium minutum*, n. gen. n. sp. dinoflagellé provocant des "eaux rouges". Vie Milieu 11: 102-105.
- Hansen, E., Eilertsen H.C., Ernstsens A. et al. 2003. Anti-mitotic activity towards sea urchin embryos in extracts from the marine haptophycean *Phaeocystis pouchetii* (Hariot) Lagerheim collected along the coast of northern Norway. Toxicon. 41: 803-812.  
[http://dx.doi.org/10.1016/S0041-0101\(03\)00034-5](http://dx.doi.org/10.1016/S0041-0101(03)00034-5)
- Houdan A., Bonnard A., Fresnel, J. et al. 2004. Toxicity of coastal coccolithophores (Prymnesiophyceae, Haptophyta). J. Plankton Res. 26: 875-883.  
<http://dx.doi.org/10.1093/plankt/fbh079>
- ICES 2011. Report of the ICES - IOC Working Group on Harmful Algal Bloom Dynamics (WGABD), 5-8 April 2011, Gothenburg, Sweden. ICES CM 2011/SSGHIE, 42 pp.
- Ichimi K., Suzuki T., Ito A. 2002. Variety of PSP toxin profiles in various culture strains of *Alexandrium tamarense* and change of toxin profile in natural *A. tamarense* population. J. Exp. Mar. Biol. Ecol. 273: 51-60.  
[http://dx.doi.org/10.1016/S0022-0981\(02\)00137-5](http://dx.doi.org/10.1016/S0022-0981(02)00137-5)
- Igarashi T., Satake M., Yasumoto T. 1996. Prymnesin-2: a potent ichthyotoxic and haemolytic glycoside isolated from the red tide algae *Prymnesium parvum*. J. Am. Chem. Soc. 118: 479-480.  
<http://dx.doi.org/10.1021/ja9534112>
- Johnsen M., Eikrem W., Olseng D.C. et al. 2010. *Prymnesium parvum*: The Norwegian Experience. J. Am. Water Resour. As. 46: 6-13.  
<http://dx.doi.org/10.1111/j.1752-1688.2009.00386.x>
- Lemée R., Pesado D., Durand-Clément M. et al. 1993. Preliminary survey of toxicity of the green alga *Caulerpa taxifolia* introduced into the Mediterranean. J. Appl. Phycol. 5: 485-493.  
<http://dx.doi.org/10.1007/BF02182507>
- Manning S.R., La Claire J.W. 2010. Prymnesins: toxic metabolites of the golden alga, *Prymnesium parvum* Carter (Haptophyta). Mar. drugs. 8: 678-704.  
<http://dx.doi.org/10.3390/md8030678>
- Martins C.A., Kulis D., Franca S. et al. 2004. The loss of PSP toxin production in a formerly toxic *Alexandrium lusitanicum* clone. Toxicon. 43: 195-205.  
<http://dx.doi.org/10.1016/j.toxicon.2003.11.023>
- Menchaca I. 2010. Desarrollo y aplicación de herramientas para la evaluación de la calidad del medio marino del País Vasco: Bioensayos con *Paracentrotus lividus* y *Corophium* sp. PhD thesis, University of the Basque Country, 220 pp.
- Moestrup Ø. 1994. Economic aspects: 'blooms', nuisance species, and toxins. In: Green, J.C. and Leadbeater, B.S.C., (eds), The Haptophyte Algae. Clarendon Press, Oxford. 265-285.
- Montoya N.G., Fulco V.K., Carignan M.O. et al. 2010. Toxin variability in cultured and natural populations of *Alexandrium tamarense* from southern South America - Evidences of diversity and environmental regulation. Toxicon. 56: 1408-1418.  
<http://dx.doi.org/10.1016/j.toxicon.2010.08.006>
- OSPAR 2005. Protocols on methods for the testing of chemicals used in the offshore oil industry. OSPAR Commission. 35 pp.
- Paul V.J., Fenical W. 1984. Bioactive terpenoids from Caribbean marine algae of the genera *Penicillus* and *Udotea* (Chlorophyta). Tetrahedron. 40: 2913-2918.  
[http://dx.doi.org/10.1016/S0040-4020\(01\)91301-8](http://dx.doi.org/10.1016/S0040-4020(01)91301-8)
- Pedrotti M.L., Marchi B., Lemée R. 1996. Effect of *Caulerpa taxifolia* secondary metabolites on the embryogenesis, larval development and metamorphosis of the sea urchin *Paracentrotus lividus*. Oceanol. Acta. 19: 255-262.
- Penna A., Fraga S., Masó M. et al. 2008. Phylogenetic relationships among the Mediterranean *Alexandrium* (Dinophyceae) species based on sequences of 5.8S gene and Internal Transcript Spacers of the rRNA operon. Eur. J. Phycol. 43: 163-178.  
<http://dx.doi.org/10.1080/09670260701783730>
- Pérez V. 2006. La utilización de anfípodos para la evaluación integrada de la calidad de sedimentos marinos y estuáricos: su aplicación en la costa vasca. PhD thesis, University of the Basque Country, 197 pp.
- Pérez S., Beiras R. 2010. The mysid *Siriella armata* as a model organism in marine ecotoxicology: comparative acute toxicity sensitivity with *Daphnia magna*. Ecotoxicology. 19: 196-206.  
<http://dx.doi.org/10.1007/s10646-009-0405-3>
- Pohnert G. 2000. Wound- activated chemical defense in unicellular planktonic algae. Angew. Chem. Int. Edit. 39: 4352-4354.
- Privitera D., Giussani V., Isola G. et al. 2012. Toxic effects of *O-*

- treopsis ovata* on larvae and juveniles of *Paracentrotus lividus*. Harmful Algae. 18: 16-23.  
<http://dx.doi.org/10.1016/j.hal.2012.03.009>
- Rommel E.J., Kohmescher N., Larson J.H. et al. 2011. Experimental study of harmful algae-zooplankton interactions and the ultimate grazing defense. Limnol. Oceanogr. 56: 461-470.  
<http://dx.doi.org/10.4319/lo.2011.56.2.0461>
- Rommel E.J., Hambright K.D. 2012. Toxin-assisted micropredation: experimental evidence shows that contact micropredation rather than exotoxicity is the role of Prymnesium toxins. Ecol. Lett. 15: 126-132.  
<http://dx.doi.org/10.1111/j.1461-0248.2011.01718.x>
- Ribo J.M., Kaiser K.L.E. 1987. *Photobacterium phosphoreum* toxicity bioassay. I. Test procedures and applications. Toxic. Assess. 2: 305-323
- Romano G., Russo G.L., Buttino I. et al. 2003. A marine diatom-derived aldehyde induces apoptosis in copepod and sea urchin embryos. J. Exp. Biol. 206: 3487-3494.  
<http://dx.doi.org/10.1242/jeb.00580>
- Sarabia R. 2002. Toxicidad y acumulación de cadmio en poblaciones de diferentes especies de *Artemia*. PhD thesis, University of Valencia, 138 pp.
- Seoane S., Eikrem W., Arluzea J. et al. 2009. Haptophytes of the Nervión River estuary, northern Spain. Bot. Mar. 52: 47-59.  
<http://dx.doi.org/10.1515/BOT.2009.027>
- Sorgeloos P., Rémiche-Van Der Wielen C., Persoone G. 1978. The use of *Artemia* nauplii for toxicity tests-A critical analysis. Ecotoxicol. Environ. Saf. 2: 249-255.
- USEPA 2001. Method for assessing the chronic toxicity of marine and estuarine sediment-associated contaminants with the amphipod *Leptocheirus plumulosus*. EPA 600/R-01/020. 102 pp.
- USEPA 2002. Short-term Methods for Estimating the Chronic Toxicity of Effluents and Receiving Waters to Marine and Estuarine Organisms. Third Edition, environmental Monitoring Systems. Laboratory, U.S. Environmental Protection Agency, Cincinnati, EPA 821/R-02/014. 464 pp.
- Varela D., Paredes J., Alves-de-Souza C. et al. 2012. Intra-regional variation among *Alexandrium catenella* (Dinophyceae) strains from southern Chile: Morphological, toxicological and genetic diversity. Harmful Algae. 15: 8-18.  
<http://dx.doi.org/10.1016/j.hal.2011.10.029>
- Yasumoto T., Oshima Y., Yamaguchi M. 1978. Occurrence of a new type of shellfish poisoning in the Tohoku district. B. Jpn. Soc. Sci. Fish. 44: 1249-1255.  
<http://dx.doi.org/10.2331/suisan.44.1249>
- Zhenxing W., Yinglin Z., Mingyuan Z. et al. 2006. Effects of toxic *Alexandrium* species on the survival and feeding rates of brine shrimp, *Artemia salina*. Acta Ecol. Sin. 26: 3942-3947.  
[http://dx.doi.org/10.1016/S1872-2032\(07\)60004-3](http://dx.doi.org/10.1016/S1872-2032(07)60004-3)
- Zon L.I., Peterson R.T. 2005. In vivo drug discovery in the zebrafish. Nat. Rev. Drug. Discov. 4: 35-44.  
<http://dx.doi.org/10.1038/nrd1606>