

## Metazoan parasites of grey mullets (Teleostea: Mugilidae) from the Mistras Lagoon (Sardinia, western Mediterranean)\*

PAOLO MERELLA and GIOVANNI GARIPPA

Dipartimento di Biologia Animale, Università degli Studi di Sassari, via Vienna, 2, 07100 Sassari, Italia.  
E-mail: malpar@uniss.it

**SUMMARY:** A total of 127 specimens of grey mullets (Teleostea: Mugilidae) from the Mistras Lagoon (Sardinia – western Mediterranean) were examined for metazoan parasites. Five species of grey mullets were found: *Chelon labrosus*, *Liza aurata*, *L. ramada*, *L. saliens* and *Mugil cephalus*. Forty-one species of parasites were identified: 14 monogeneans, 5 crustaceans, 18 digeneans, 3 nematodes and 1 acanthocephalan; 26 of them being reported from Sardinia for the first time. Findings of faunal interest are: *Ergenstrema* sp. (Monogenea) in *L. saliens*; *Caligus pageti* (Crustacea) in *L. aurata* and *L. saliens*; *Saturnius* sp. (Digenea) in *C. labrosus*, *L. aurata* and *L. saliens*; *Capillaria* sp. (Nematoda) in *C. labrosus*, *L. aurata* and *M. cephalus*. The dynamics and distribution of ectoparasites were partially influenced by salinity, while the interpretation of the results for endoparasites was more complex. The data obtained show the richness of the parasite fauna of grey mullets from the Mistras Lagoon, providing additional environmental information on this body of water.

**Key words:** ectoparasites, endoparasites, Teleostea, Mugilidae, western Mediterranean, Sardinia.

### INTRODUCTION

Grey mullets (family Mugilidae) are a cosmopolitan family of teleost fishes. Extensive and semi-intensive exploitation of these fishes has been practised for centuries in the Mediterranean region, south-east Asia, Taiwan, Japan and Hawaii, in lagoons and man-made ponds (Nash and Koningsberger, 1981). In the brackish waters of the western Mediterranean this family is represented by 5 species (Tortonese, 1975; Bauchot, 1987): *Chelon labrosus* (Risso, 1826), *Liza aurata* (Risso, 1810), *L. ramada* (Risso, 1826), *L. saliens* (Risso, 1810) and *Mugil cephalus* Linnaeus, 1758.

\*Received November 3, 2000. Accepted January 22, 2001.

A comprehensive review of parasites of grey mullets world-wide has been published by Paperna and Overstreet (1981). Many data about the systematics of the parasites of mugilids are available in scientific papers, but works dealing with quantitative data and ecology are scarce; some research has been carried out in the United States (Rawson, 1973, 1976; Skinner, 1975), the Black Sea (Reshetnikova, 1955) and the Mediterranean (Ben Hassine, 1983; Radujkovic and Raibaut, 1989a; D'Amelio *et al.*, 1996; D'Amelio and Gerasi, 1997).

In the present paper the metazoan parasite fauna of grey mullets from the Mistras Lagoon (Sardinia, western Mediterranean) (Fig. 1) is described. The investigation was primarily concerned with parasites of the skin, gills and digestive tract. Most results of




FIG. 1. – The Mistras Lagoon, site of study.

studies in the same area only report the occurrence, without providing quantitative data on infestation levels (Arru *et al.*, 1988, 1995; Garippa *et al.*, 1988; Sanna and Garippa, 1989; Merella and Garippa, 1998, 2000), so this research may be considered a first contribution to an understanding of the ecology of the parasites of mugilids from Sardinia.

## MATERIALS AND METHODS

The Mistras Lagoon (Fig. 1) is a brackish body of water of about 450 ha. The salinity is relatively high (30‰–40‰), due to the large entrance channel and to the limited inputs of fresh water; the average temperature is about 20°C, with values ranging from 10°C to 30°C.

Fishes were sampled in the fish barrier of the Mistras Lagoon, where this family is extensively exploited because of the natural recruitment of grey mullets in Mediterranean lagoons (Chauvet, 1988; Merella, 2000). Sampling was carried out in November and December 1997 and from May to August 1998; in January–April 1998 it was not possible to take samples due to lack of fish because of the low temperature. A total of 127 grey mullets belonging to the species *Chelon labrosus*, *Liza aurata*, *L. ramada*, *L. saliens* and *Mugil cephalus* were collected. The number of each species examined and the range of total length (TL) are given in Table 1.

Each fish was examined for skin parasites, then the gills and the digestive tract were extracted, stored in single bags and frozen at -20°C. Subsequently, the samples were placed in Petri dishes and examined under a stereoscope for parasites; all the specimens were counted and stored in 70% ethyl

TABLE 1. – Number and range of total length (TL) of the host species examined

Species	Number	Range of TL (cm)
<i>Chelon labrosus</i>	24	16-37
<i>Liza aurata</i>	43	21-32
<i>Liza ramada</i>	11	22-31
<i>Liza saliens</i>	38	18-30
<i>Mugil cephalus</i>	11	22-40

alcohol. For identification, monogeneans and digenleans were mounted in glycerine-jelly, digenleans and polyopisthocotylean monogeneans were pre-stained with borax carmine; crustaceans, acanthocephalans and nematodes were cleared with lactophenol and examined as temporary mounts. Illustrations of each parasite species were made with the aid of a camera lucida; drawings of specimens which could not be classified to species or genus level were given an identification code.

Among the papers consulted for the identification of parasite species were: Yamaguti (1963b), Euzet and Combes (1969), Euzet and Suriano (1977), Anderson (1981a) and Euzet and Sanfilippo (1983), for monogeneans; Yamaguti (1958), Farès and Maillard (1974), Overstreet (1977), Ramadan *et al.* (1989) and Dimitrov *et al.* (1998), for digenleans; Yamaguti (1961), Moravec (1982) and Orechia and Paggi (1987), for nematodes; Petrotchenko (1956) and Yamaguti (1963c), for acanthocephalan; Russel (1925), Brian (1935), Yamaguti (1963a), Kabata (1979, 1992) and Ben Hassine (1983), for copepods; Trilles (1975) and Charti-Cheikhrouha *et al.* (2000), for isopods. Radujkovic and Raibaut monograph (1989a) was particularly useful for identifying almost all the parasitic groups.

TABLE 2. – Parasite-host list (Prev.=prevalence; Range Int. = range of intensity; M.Int. = mean intensity; Abund.= abundance; S.D. = standard deviation. Unidentified digeneans are listed by codes beginning with “dig”)

Taxa - Species		ECTOPARASITES			
	Host	Prev.	Range Int.	M.Int.±S.D.	Abund.±S.D.
<b>Monogenea</b>					
<i>Ligophorus acuminatus</i> Euzet et Suriano, 1977	<i>L. saliens</i>	100%	2-565	159.74±142.24	159.74±142.24
<i>Ligophorus angustus</i> Euzet et Suriano, 1977	<i>C. labrosus</i>	75%	2-114	27.89±33.14	20.92±31.04
<i>Ligophorus chabaudi</i> Euzet et Suriano, 1977	<i>M. cephalus</i>	100%	3-37	15.09±10.39	15.09±10.39
<i>Ligophorus confusus</i> Euzet et Suriano, 1977	<i>L. ramada</i>	64%	4-34	16.14±13.37	10.27±13.18
<i>Ligophorus heteronchus</i> Euzet et Suriano, 1977	<i>L. saliens</i>	97%	4-563	104.78±114.61	102.03±114.32
<i>Ligophorus imitans</i> Euzet et Suriano, 1977	<i>L. ramada</i>	91%	1-39	13.10±12.42	11.91±12.43
<i>Ligophorus macrocolpos</i> Euzet et Suriano, 1977	<i>L. saliens</i>	95%	9-223	58.25±50.15	55.18±50.53
<i>Ligophorus minimus</i> Euzet et Suriano, 1977	<i>L. saliens</i>	47%	2-67	19.67±17.91	9.32±15.70
<i>Ligophorus szidati</i> Euzet et Suriano, 1977	<i>L. aurata</i>	63%	1-197	32.37±49.70	20.33±42.19
<i>Ligophorus vanbenedenii</i> (Parona et Perugia, 1890)	<i>L. aurata</i>	98%	3-141	38.67±35.31	37.77±35.38
<i>Ergenstrema labrosi</i> Anderson, 1981	<i>C. labrosus</i>	21%	2-5	2.80±1.30	0.58±1.28
<i>Ergenstrema mugilis</i> Paperna, 1964	<i>L. ramada</i>	36%	2-4	2.75±0.96	1.00±1.48
<i>Ergenstrema</i> sp.	<i>L. saliens</i>	8%	4-26	13.33±11.37	1.05±4.50
<i>Solostamenides mugilis</i> (Vogt, 1878)	<i>C. labrosus</i>	4%	1	1.00	0.04±0.20
<b>Crustacea</b>					
<i>Nipergasilus bora</i> (Yamaguti, 1939)	<i>C. labrosus</i>	33%	2-5	2.88±1.13	0.96±1.52
“	<i>L. saliens</i>	16%	1-3	1.83±0.75	0.29±0.73
“	<i>M. cephalus</i>	27%	1-2	1.33±0.58	0.36±0.67
<i>Caligus apodus</i> (Brian, 1924)	<i>C. labrosus</i>	63%	1-8	2.53±2.00	1.58±2.00
“	<i>L. ramada</i>	64%	1-4	2.00±1.15	1.27±1.35
“	<i>L. saliens</i>	37%	1-6	2.29±1.44	0.84±1.41
“	<i>M. cephalus</i>	55%	1-2	1.17±0.41	0.64±0.67
<i>Caligus mugilis</i> Brian, 1935	<i>C. labrosus</i>	4%	1	1.00	0.04±0.200
<i>Caligus pageti</i> Russell, 1925	<i>L. aurata</i>	2%	1	1.00	0.02±0.15
<i>L. saliens</i>	<i>L. saliens</i>	3%	1	1.00	0.03±0.16
<i>Nerocila orbignyi</i> (Guérin-Méneville, 1832)	<i>L. aurata</i>	2%	1	1.00	0.02±0.15
<b>ENDOPARASITES</b>					
<b>Digenea</b>					
<i>Dicrogaster contractus</i> Looss, 1902	<i>C. labrosus</i>	54%	4-230	62.38±77.27	33.79±64.21
“	<i>L. aurata</i>	70%	1-1160	135.40±219.73	94.47±193.12
“	<i>L. ramada</i>	55%	2-40	11.67±14.40	6.36±11.87
“	<i>L. saliens</i>	87%	6-3208	265.12±586.55	230.24±552.99
<i>Haploporus benedeni</i> (Stossich, 1887)	<i>C. labrosus</i>	67%	3-737	290.81±232.76	193.88±234.40
“	<i>L. ramada</i>	36%	1-34	17.25±16.66	6.27±12.61
“	<i>L. saliens</i>	11%	1-45	16.25±19.62	1.71±7.53
<i>Lecithobotrys putrescens</i> Looss, 1902	<i>L. saliens</i>	34%	1-25	9.62±7.54	3.29±6.31
<i>Saccocoelium obesum</i> Looss, 1902	<i>C. labrosus</i>	13%	5-45	19.33±22.28	2.42±9.26
“	<i>L. aurata</i>	81%	1-47	13.31±13.28	10.84±13.05
“	<i>L. ramada</i>	9%	8	8	0.73±2.41
“	<i>L. saliens</i>	66%	1-45	7.24±9.26	4.76±8.23
<i>Saccocoelium tensum</i> Looss, 1902	<i>C. labrosus</i>	50%	6-152	44.33±40.02	22.17±35.76
“	<i>L. aurata</i>	72%	1-114	16.71±23.42	12.05±21.20
“	<i>L. ramada</i>	100%	4-70	19.36±20.21	19.36±20.21
“	<i>L. saliens</i>	5%	2	2.00±0.00	0.11±0.45
<i>Haplosplanchnus pachysomus</i> (Eysenhardt, 1829)	<i>L. ramada</i>	9%	1	1.00	0.09±0.30
“	<i>L. saliens</i>	16%	1-2	1.33±0.52	0.21±0.53
“	<i>M. cephalus</i>	82%	1-26	8.89±9.47	7.27±9.20
<i>Schikhobalotrema</i> sp.	<i>C. labrosus</i>	25%	4-213	111.50±86.07	27.88±63.58
“	<i>L. ramada</i>	9%	1	1.00	0.09±0.30
<i>Saturnius papernai</i> Overstreet, 1977	<i>M. cephalus</i>	91%	1-26	12.80±8.94	11.64±9.32
<i>Saturnius</i> sp.	<i>C. labrosus</i>	88%	4-310	58.52±71.12	51.21±69.20
“	<i>L. aurata</i>	42%	1-35	8.00±9.77	3.35±7.39
“	<i>L. ramada</i>	36%	1-6	4.00±2.16	1.45±2.34
“	<i>L. saliens</i>	24%	1-62	10.11±19.54	2.39±10.08
digcl1	<i>C. labrosus</i>	4%	43	43.00	1.79±8.78
digla1	<i>L. aurata</i>	12%	1-8	4.00±2.55	0.47±1.52
digls1	<i>L. saliens</i>	8%	2-40	17.33±20.03	1.37±6.64
digls2	<i>L. saliens</i>	29%	1-101	31.91±37.07	9.24±24.22
digmc1	<i>M. cephalus</i>	100%	9-2170	628.64±714.24	628.64±714.24
digmc2	<i>M. cephalus</i>	55%	1-78	22.00±30.94	12±24.71
digmc3	<i>M. cephalus</i>	45%	4-8	5.20±1.64	2.36±2.91
digmc4	<i>M. cephalus</i>	9%	8	8.00	0.73±2.41
digmc5	<i>M. cephalus</i>	9%	33	33.00	3.00±9.95
<b>Nematoda</b>					
<i>Capillaria</i> sp.	<i>C. labrosus</i>	4%	1	1.00	0.04±0.20
“	<i>L. aurata</i>	2%	1	1.00	0.02±0.15
“	<i>M. cephalus</i>	27%	1-3	2.00±1.00	0.55±1.04
<i>Contracaecum</i> sp.	<i>L. aurata</i>	9%	1-2	1.25±0.50	0.12±0.39
<i>Cucullanus bioccai</i> Orecchia et Paggi, 1987	<i>M. cephalus</i>	55%	1-4	1.67±1.21	0.91±1.22
<b>Acanthocephala</b>					
<i>Neoechinorhynchus agilis</i> (Rudolphi, 1819)	<i>C. labrosus</i>	54%	1-35	10.85±11.96	5.88±10.25
“	<i>L. aurata</i>	16%	1-6	3.00±2.16	0.49±1.39
“	<i>L. ramada</i>	18%	1	1.00±0.00	0.18±0.40
“	<i>L. saliens</i>	32%	1-5	2.92±1.56	0.92±1.62
“	<i>M. cephalus</i>	91%	1-25	5.60±7.52	5.09±7.33

The infection level of each parasite species was analysed by calculating prevalence, range of intensity, mean intensity and abundance (Margolis *et al.*, 1982).

## RESULTS

Parasites were found in 100% of the specimens examined. Table 2 summarises the parasite-host list with indications of prevalence, range of intensity, mean intensity ( $\pm$ standard deviation) and abundance ( $\pm$ standard deviation). A total of 41 species were identified: 19 ectoparasites and 22 endoparasites. All these specimens were adult stages, except some copepods of the genus *Caligus* and the nematodes belonging to the genus *Contracaecum*.

Among the ectoparasites, the genus *Ligophorus* was the most frequent and abundant, with prevalence ranging from 47% to 100% and intensity up to 565. Other ancyrocephalids, *Ergenstrema* spp. were fairly common, but not very abundant. *Caligus apodus* and *Nipergasilus bora* were the most prevalent crustaceans. Of the endoparasites, adult digeneans were most represented, and within this taxon haploporids were the most frequent and abundant family. In addition, other families, such as haplosplanchnids and hemiurids, were frequently found and abundant. Within the digeneans, nine taxa could not be classified to species or genus level because their morphometric characteristics did not seem to coincide with those of other existing species, and they are reported in Table 2 with an identification code (digcl1, digla1, digls1, digls2, digmc1, digmc2, digmc3, digmc4, digmc5). The nematodes *Capillaria* sp. and *Cucullanus bioccai* were well represented in *Mugil cephalus* and, among the acanthocephalans, *Neoechinorhynchus agilis* had relatively high prevalence in all 5 host species.

## DISCUSSION

The results provided new data about the parasites of mugilids. Of the 41 species found, 26 are new records for Sardinia: 14 monogeneans (*Ligophorus* spp., *Ergenstrema* spp. and *Solostamenides mugilis*), 4 crustaceans (*Nipergasilus bora*, *Caligus mugilis*, *C. pageti* and *Nerocila orbignyi*), 4 digenans (*Lecithobotrys putrescens*, *Saccocoelium tensum*, *Saturnius papernai*, *Saturnius* sp.), 3 nematodes (*Capillaria* sp., *Contracaecum* sp. and *Cucullanus bioccai*) and 1 acanthocephalan (*Neoechi-*

*norhynchus agilis*). In addition, 9 taxa of digeneans were not identified. All these species must be added to the others formerly recorded in the island (Arru *et al.*, 1988, 1995; Garippa *et al.*, 1988; Sanna and Garippa, 1989), 3 of which were not found in this study: *Metamicrocotyla cephalus* (Azim, 1939), *Ergasilus lizae* Krøyer, 1863 and *Floridosentis* sp.

Records of faunal interest, possibly findings of new species or new hosts, or else first reports of a parasite species for Italian marine fauna, are: *Ergenstrema* sp. in *Liza saliens*; *C. pageti* in *Liza aurata* and *L. saliens*; *Saturnius* sp. in *Chelon labrosus*, *L. aurata* and *L. saliens*; *Capillaria* sp. in *C. labrosus*, *L. aurata* and *Mugil cephalus*.

Parasites of the genus *Ergenstrema* have never been recorded in *L. saliens*. The morphometric characteristics of the specimens found do not match those of the congeners parasitising *C. labrosus* and in *Liza ramada*, particularly in the length and coiling of the vagina (Anderson, 1981a; Radujkovic and Euzet, 1989), and they probably represent a new species.

*C. pageti* is a new record for Italian marine fauna. This species has been found in Algeria, Tunisia and France (Ben Hassine, 1983). Brian (1935), after examining numerous mugilids from Italy, affirmed that *C. pageti* was present only in the eastern and southern Mediterranean.

Species of the genus *Saturnius* were formerly recorded only in *M. cephalus* (Overstreet, 1977; Dimitrov *et al.*, 1998) and *L. ramada* (D'Amelio *et al.*, 1996; Dimitrov *et al.*, 1998). Overstreet (1977), who found *Saturnius* species only in *M. cephalus*, concluded that if new hosts for this genus exist, the species would have to be a new *Saturnius* one, due to the specialisation of this genus.

This is the first record of adult stages of capillariids in the digestive tract of mugilids from the Mediterranean; formerly, only Rawson (1973) and Moravec *et al.* (1999) found nematodes of this family in mugilids from Georgia and Mexico respectively, apart from the larval stages recorded in the Philippines (Paperna and Overstreet, 1981). Because of problems with identification of the fragmented specimens, and the unsatisfactory status of the taxonomy of capillariids (Moravec, 1982; Moravec *et al.*, 1999), the individuals belonging to this family were generically assigned to the genus *Capillaria*.

The prevalence values of *Ligophorus* spp. were higher than those obtained in the Boka Kotorska Bay (Montenegro) from the same hosts, ranging from 28% to 83% (Radujkovic and Euzet, 1989) vs. 47%-100% (Table 2); while the prevalence values of

*Ergenstrema* spp. were lower, ranging 21% to 36% (Table 2) vs. 46% to 57% (Radujkovic and Euzet, 1989). These differences could be explained by the salinity of the lagoon, higher in the site of the study (30‰-40‰) than in the Boka Kotorska Bay, ranging from 1‰ to 37‰ (Stjepcevic, 1974). This may be a limiting factor for *Ergenstrema* and favour the *Ligophorus* species (Anderson, 1981b).

Prevalences of *N. bora* were similar to those obtained in the Boka Kotorska Bay by Radujkovic and Raibaut (1989b), while *C. apodus* showed higher prevalence values than those found by the same authors: 2%-3% vs. 37%-64% (Table 2), confirming the preference of this species for the brackish environments (Ben Hassine, 1983).

The only skin ectoparasite found was 1 specimen of *N. orbignyi*. The rarity of this species in the Mistras Lagoon is probably due to the high salinity. In the adjacent Cabras Lagoon, characterised by oligohaline waters, the prevalence and intensity of *N. orbignyi* is much higher (P. Merella, unpublished data).

Prevalence and mean intensity of digeneans were mostly higher than those reported from the Boka Kotorska Bay by Radujkovic *et al.* (1989). These values were also higher for *Contracaecum* sp. in *L. aurata* compared to the results of Rawson (1973) and Petter and Radujkovic (1989); while the prevalence of *Capillaria* sp. in *M. cephalus* was close to that reported by Rawson (1973).

The prevalence values of *N. agilis* were elevated in all 5 species of hosts, and higher than those indicated for other Mediterranean regions by Radujkovic (1989) and D'Amelio *et al.* (1996): 1%-30% vs. 16%-91% (Table 2).

Although fish were sampled during 6 months of the year, their number in each monthly sample was too low for any conclusions as to their possible seasonal patterns. However, ectoparasite distributions were partially influenced by salinity, while the interpretation of the results regarding endoparasites was more complex, because of numerous biotic and abiotic factors (Rawson, 1973).

Concerning the gill parasites of *C. labrosus*, it was observed that the prevalence of *L. angustus* was higher in July-August, when the salinity was about 40‰, while *E. labrosi* peaked in June, when salinity fell to about 30‰, supporting the hypothesis of Anderson (1981b) that *L. angustus* prefers higher salinities compared to *E. labrosi*. Analogously, *C. pageti* was found on *L. aurata* and *L. saliens* only in November, when salinity was about 40‰, confirming the preference of this species for hyperhaline

waters (Ben Hassine, 1983). One single specimen of *N. orbignyi* appeared in June, when salinity was about 30‰, while, as mentioned above, in the adjacent Cabras Lagoon, characterised by oligohaline waters, the prevalence and intensity of this species was notably higher all the year round (P. Merella, unpublished data).

The results obtained show the richness of the parasite fauna of grey mullets from the Mistras Lagoon; and although the parasites of fish cannot at present be considered as reliable bioindicators, they provide useful additional environmental information on the body of water studied (D'Amelio and Gerasi, 1997; Kennedy, 1997; Overstreet, 1997).

The numerous new records for the mugilid parasite fauna, the possible finding of new parasite species and the difficulties encountered in the identification of some species, particularly digeneans, point to the need for further investigations, to extend knowledge of the biology and ecology of metazoan parasites of grey mullets.

## ACKNOWLEDGEMENTS

We are grateful to: Prof. Lia Paggi and her research staff (Istituto di Parassitologia, Università degli studi di Roma "La Sapienza") for their useful suggestions; Dr. Fulvio Salati and his working group (Istituto Zooprofilattico Sperimentale della Sardegna "G. Pegreff") for their collaboration; and Dr. Tomáš Scholz (Institute of Parasitology ASCR - Ceske Budejovice) for his helpfulness and the critical reading of the paper. Particular thanks are due to Mr. Salvatore Manca and all the fisherman of the Mistras Lagoon for their courtesy and help during the sampling. Mrs. Vera Walker revised the English version of the manuscript. This research was funded by MURST (60%).

## REFERENCES

- Anderson, M. – 1981a. *Ergenstrema labrosi* sp. nov. (Monogenea) on the gills of the thick-lipped grey mullet *Chelon labrosus* at Plymouth. *J. Mar. Biol. Ass. UK.*, 61: 827-832.  
Anderson, M. – 1981b. The change with host age of the composition of the ancyrocephaline (Monogenean) populations of parasites on thick-lipped grey mullets at Plymouth. *J. Mar. Biol. Ass. UK.*, 61: 833-842.  
Arru, E., A. Leoni and G. Garippa. – 1988. Contributo alla conoscenza della parassitofauna dei pesci delle acque costiere ed interne della Sardegna. *Parassitologia*, 30: 14-16.  
Arru, E., G. Garippa and M.L. Sanna. – 1995. I parassiti dei Mugilidi delle acque salmastre della Sardegna. *Biol. Mar. Medit.*, 2: 571-572.  
Bauchot, M.-L. – 1987. Poissons osseux. In: W. Fischer, M.-L.

- Bauchot & M. Schneider (eds). *Fiches FAO d'identification des espèces pour les besoins de la pêche. (Révision 1). Méditerranée et Mer Noire. Zone de pêche 37. Vol. II. Vertébrés*, pp. 891-1422. FAO, Rome.
- Ben Hassine, O.K. – 1983. *Les Copépodes parasites de Poissons Mugilidae en Méditerranée (côtes françaises et tunisiennes) – Morphologie, Bio-écologie, Cycles évolutifs*. Ph. D. thesis, Univ. Sciences et Techniques du Languedoc, Montpellier.
- Brian, A. – 1935. I *Caligus* parassiti dei pesci del Mediterraneo (Copepodi). *Ann. Mus. Civ. Storia Nat. Genova*, 57: 152-211.
- Charti-Cheikhrouha, F., W. Zghidi, L. Ould Yarba and J.P. Trilles. – 2000. Les Cymothoidae (Isopodes parasites de poissons) des côtes tunisiennes: écologie et indices parasitologiques. *Syst. Parasitol.*, 46:143-150.
- Chauvet, C. – 1988. Manuel sur l'aménagement des pêches dans les lagunes côtières: la bodrigue méditerranéenne. *Fao Doc. Tech. Pêches*.
- D'Amelio, S., D. Di Cave, P. Orecchia and L. Paggi. – 1996. Le comunità dei metazoi parassiti di *Liza ramada* (Risso, 1826) (Pisces: Mugilidae) nei laghi di Sabaudia (LT) e del Fusaro (NA). *Suppl. Ric. Biol. Selvaggina*, 24: 671-676.
- D'Amelio, S. and L. Gerasi. – 1997. Evaluation of environmental deterioration by analysing fish parasites biodiversity and community structure. *Parasitol.*, 39: 237-241.
- Dimitrov, G.I., F.K. Arapoglou and B.B. Georgiev. – 1998. First record of *Saturnius papernai* Overstreet, 1977 in the Black Sea, with a review of the genus *Saturnius* Manter, 1969 (Digenea, Bunocotylidae). *Syst. Parasitol.*, 40: 43-48.
- Euzet, L. and C. Combes. – 1969. Contribution à l'étude des Microcotylidae (Monogenea) Parasites de *Mugil cephalus* L. (Teleostei). *Parazit. sbornik*, 24: 91-105.
- Euzet, L. and M.D. Suriano. – 1977. *Ligophorus* n. g. (Monogenea: Ancyrocephalidae) parasite des Mugilidae (Téléostéens) en Méditerranée. *Bull. Mus. natn. Hist. Nat., Paris*, 3<sup>e</sup> sér., 472, Zool., 329: 799-822.
- Euzet, L. and D. Sanfilippo. – 1983. *Ligophorus parvicirrus* n. sp. (Monogenea, Ancyrocephalidae) parasite de *Liza ramada* (Risso, 1826) (Teleostei, Mugilidae). *Ann. Parasitol. Hum. Comp.*, 1983: 325-335.
- Farès, A. and C. Maillard. – 1974. Recherches sur quelques Haploporeidae (Trematoda) parasites des Muges de Méditerranée Occidentale: systématique et cycles évolutifs. *Z. Parasitenk.*, 45: 11-43.
- Garippa, G., E. Sanna and G.F. Cuzzupoli. – 1988. Segnalazione di *Haplosplanchnus pachysomus* in *Mugil cephalus* dello Stagno di "Is Benas" (Prov. Di Oristano). *Parassitologia*, 30: 77-78.
- Kabata, Z. – 1979. *Parasitic Copepoda of British Fishes*. The Ray Society, London.
- Kabata, Z. – 1992. *Copepods Parasitic on Fishes*. Synopses of the British Fauna (New Series), 47. The Linnean Society of London & The Estuarine and Coastal Sciences Association, London.
- Kennedy, C.R. – 1997. Freshwater fish parasites and environmental quality: an overview and caution. *Parassitologia*, 39: 249-254.
- Margolis, L., G.W. Esch, J.C. Holmes, A.M. Kuris and G.A. Schad. – 1982. The use of ecological terms in parasitology (Report of an ad hoc committee of the American Society of Parasitologists). *J. Parasitol.*, 68: 131-133.
- Merella, P. – 2000. Per le peschiere degli stagni del Sinis. *Civiltà del Mare*, 11(1): 4-6.
- Merella, P. and G. Garippa. – 1998. Parasites of grey mullets from the Oristano lagoons (Sardinia). *Parassitologia*, 40(Suppl 1): 106.
- Merella, P. and G. Garippa. – 2000. Parasite fauna of mugilids (Osteichthyes: Mugilidae) from the Mistra Lagoon (Sardinia – western Mediterranean). *Parassitologia*, 42(Suppl 1): 177.
- Moravec, F. – 1982. Proposal of a new systematic arrangement of nematodes of the family Capillariidae. *Folia Parasit.*, 29: 119-132.
- Moravec, F., G. Salgado-Maldonado and J. Caspeta-Mandujano. – 1999. Capillariids (Nematoda, Capillariidae) from *Agonostomus monticola* and *Gobiomorus* spp. (Pisces, Mugilidae and Eleotridae) from fresh waters in Mexico. *Acta Parasit.*, 44: 180-187.
- Nash, C.E. and R.M. Koningsberger. – 1981. Artificial propagation. In: O.H. Oren (ed.), *Aquaculture of grey mullets*, pp. 265-312. Cambridge University Press, Cambridge.
- Orecchia, P. and L. Paggi. – 1987. *Cucullanus bioccaii* n. sp. (Nematoda: Cucullanida) parassita di *Mugil cephalus* L. *Parasitologia*, 29: 71-74.
- Overstreet, R.M. – 1977. A revision of *Saturnius* Manter, 1969 (Hemiuroidae: Bunocotylinae) with description of two new species from the striped mullet. In: R. Lamothe-Argumedo et al. (eds.), *Excerpta Parasitologica en Memoria del Doctor Eduardo Caballero y Caballero*, pp. 273-284. Universidad Nacional Autónoma de México, México City.
- Overstreet, R.M. – 1997. Parasitological data as monitors of environmental health. *Parassitologia*, 39: 169-175.
- Paperna, I. and R.M. Overstreet. – 1981. Parasites and diseases of mullets (Mugilidae). In: O.H. Oren (ed.), *Aquaculture of grey mullets*, pp. 411-493. Cambridge University Press, Cambridge.
- Petrotschenko, V.I. – 1956. *Acanthocephala of domestic and wild animals*. Vol. I. Moskow. Israel Progr. Sci. Transl. Jerusalem.
- Petter, A.J. and B.M. Radujkovic. – 1989. Parasites des poissons marins du Monténégro: Nematodes. *Acta Adriat.*, 30: 195-236.
- Radujkovic, B.M. – 1989. Parasites des poissons marins du Monténégro: Acanthocephales. *Acta Adriat.*, 30: 189-194.
- Radujkovic, B.M. and L. Euzet. – 1989. Parasites des poissons marins du Monténégro: Monogènes. *Acta Adriat.*, 30: 51-135.
- Radujkovic, B.M., P. Orecchia and L. Paggi. – 1989. Parasites des poissons marins du Monténégro: Digenes. *Acta Adriat.*, 30: 137-187.
- Radujkovic, B.M. and A. Raibaut. – 1989a. Faune des parasites des poissons marins des côtes du Monténégro (Adriatic Sud). *Acta Adriat.*, 30: 1-324.
- Radujkovic, B.M. and A. Raibaut. – 1989b. Parasites des poissons marins du Monténégro: Copépodes. *Acta Adriat.*, 30: 237-278.
- Ramadan, M.M., M.F.A. Saoud, A.A. Ashour and D.A. Mansour. – 1989. A review of the genus *Saccocoelium* Looss, 1902 (Trematoda, Haploporidae) with a redescription of *S. tensum* Looss, 1902 and a description of *S. gohari* sp. n. from Egyptian Lake Qarun fishes. *Acta Parasitologica Polonica*, 34: 125-135.
- Rawson, M.V. Jr. – 1973. The development and seasonal abundance of the parasites in stripped mullet *Mugil cephalus* L. and mummichogs *Fundulus heteroclitus* (L.). Dissertation, University of Georgia, Athens.
- Rawson, M.V. Jr. – 1976. Population biology of parasites of stripped mullet *Mugil cephalus* L. I - Monogenea. *J. Fish Biol.*, 9: 185-194.
- Reshetnikova, A.V. – 1955. Parasite fauna of *Mugil cephalus* in the Black Sea. *Trudy Karadah's koi Biologichnoi Stantsii*, 13: 71-95.
- Russel, F.S. – 1925. A new species of *Caligus* from Egypt, *Caligus pageti*, sp. n. *Ann. Mag. Nat. Hist.*, 15 (9): 611-618.
- Sanna, E. and G. Garippa. – 1989. *Saccocoelium, Haploporus e Dicrogaster* spp. in *Mugil cephalus* della Sardegna. *Boll. Soc. It. Patol. Ittica*, 2: 6-10.
- Skinner, R. – 1975. Parasites of the stripped mullet, *Mugil cephalus*, from Biscayne bay, Florida, with descriptions of a new genus and three new species of trematodes. *Bull. Mar. Sci.*, 5: 318-345.
- Stjepcevic, J. – 1974. Ekologija dagnje (*Mytilus galloprovincialis* Lamk.) i kamenice (*Ostrea edulis* L.) u gajilištima Bokokotorskih zaliva – Ecology of mussel (*Mytilus galloprovincialis* Lamk.) and oyster (*Ostrea edulis* L.) in cultures of Boka Kotorska Bay. *Stadia Marina*, 7: 5-164.
- Tortonese, E. – 1975. *Fauna d'Italia, Osteichthyes* - Vol. XI. Calderini, Bologna.
- Trilles, J.-P. – 1975. Les Cymothoidae (Isopoda, Flabellifera) des côtes françaises II. Les Anilocridae Schioedte et Meinert, 1881. Genres *Anilocra* Leach, 1818 et *Nerocila* Leach, 1818. *Bull. Mus. natn. Hist. nat., Paris*, 3<sup>e</sup> sér., 290, zool., 200: 347-378.
- Yamaguti, S. – 1958. *Systema Helmintum - Volume I - The Digenetic Trematodes of Vertebrates, Part I, II*. Interscience Publishers, New York.
- Yamaguti, S. – 1961. *Systema Helmintum - Volume III - Nematodes*. Interscience Publishers, New York.
- Yamaguti, S. – 1963a. *Parasitic Copepoda and Branchiura of fishes*. Interscience Publishers, New York.
- Yamaguti, S. – 1963b. *Systema Helmintum - Volume IV - Monogenea and Aspidocotylea*. Interscience Publishers, New York.
- Yamaguti, S. – 1963c. *Systema Helmintum - Volume V - Acanthocephala*. Interscience Publishers, New York.

Scient. ed.: P. Oliver